



Impact of trifluridine/tipiracil plus bevacizumab on tumor shrinkage and depth of response in refractory metastatic colorectal cancer: analysis of the SUNLIGHT trial

Julien Taieb^{a,b,*}, Dominik P. Modest^{c,d}, Marwan Fakih^e, Fortunato Ciardiello^f, Eric Van Cutsem^g, Elena Elez^h, Cristina Gravalosⁱ, Arinilda Campos Bragagnoli^j, Valentine Barbou^k, Lucas Roby^k, Nadia Amellal^k, Gerald W. Prager^l

^a Department of Gastroenterology and Gastrointestinal Oncology, CARPEM Comprehensive Cancer Center, Georges-Pompidou European Hospital, AP-HP, Université Paris-Cité, Paris, France

^b Centre de Recherche des Cordeliers, INSERM, CNRS, Université Paris-Cité, Sorbonne Université, USPC, Equipe Labellisée Ligue Nationale Contre le Cancer, SIRIC CARPEM, Paris, France

^c Department of Hematology, Oncology and Cancer Immunology (CVK), Charité-Universitätsmedizin, Berlin, Germany

^d German Cancer Consortium (DKTK), partner site Berlin, German Cancer Research Centre (DKFZ), Heidelberg, Germany

^e City of Hope Comprehensive Cancer Center, Duarte, USA

^f Università degli Studi della Campania Luigi Vanvitelli, Naples, Italy

^g University Hospitals Gasthuisberg and University of Leuven (KUL), Leuven, Belgium

^h Vall d'Hebron Hospital Campus and Institute of Oncology (VHIO), UVic-UCC, Barcelona, Spain

ⁱ Medical Oncology Department, 12 de Octubre University Hospital, Madrid, Spain

^j Fundacao Pio XII Hospital Cancer De Barretos, Barretos, Brazil

^k Servier International Research Institute, Suresnes, France

^l Medical University Vienna, Vienna, Austria

ARTICLE INFO

Keywords:

Colorectal cancer
Trifluridine/tipiracil
Bevacizumab
Response parameters
Depth of response

ABSTRACT

Objectives: This *post hoc* analysis of the SUNLIGHT trial sought to assess the response to treatment with trifluridine/tipiracil (FTD/TPI) + bevacizumab and FTD/TPI in patients with refractory metastatic colorectal cancer using tumor shrinkage (TS), early TS (ETS), duration of TS (DTS) and depth of response (DpR) as response-related parameters.

Methods: TS was defined as any decrease from baseline of the sum of the longest diameter of target lesions. TS at first assessment was specified as ETS. DpR was defined as the maximum percentage change from baseline of the sum of the longest diameters of target lesions. DTS was defined as the time from first TS to first increase in tumor size, progressive disease, or death.

Results: In the FTD/TPI + bevacizumab group, 48 % had TS and 39 % had ETS. In the FTD/TPI group, 21 % had TS and 17 % had ETS. In patients achieving ETS, median DTS was prolonged with FTD/TPI + bevacizumab compared to FTD/TPI (3.8 versus 2.1 months; HR: 0.34 [95 % CI: 0.22, 0.53]; $P < 0.0001$). Magnitude of DpR was greater with FTD/TPI + bevacizumab than with FTD/TPI.

Conclusion: The survival benefit of treatment with FTD/TPI + bevacizumab versus FTD/TPI is likely associated with the improvement of ETS and DpR.

1. Introduction

The phase 3 SUNLIGHT trial demonstrated the benefit of treatment with trifluridine/tipiracil (FTD/TPI) + bevacizumab on overall survival

(OS) and progression-free survival (PFS) compared to FTD/TPI in patients with metastatic colorectal cancer (mCRC) who had received no more than two lines of treatment and had progressed, or were intolerant to, their last line of treatment [1]. SUNLIGHT predominately recruited

* Correspondence to: Department of Gastroenterology and Digestive Oncology, European Georges Pompidou Hospital, AP-HP and Paris-Cité University, SIRIC CARPEM Comprehensive Cancer Center, 20 Rue Leblanc, Paris 75015, France.

E-mail address: jtaieb75@gmail.com (J. Taieb).

<https://doi.org/10.1016/j.ejca.2025.115644>

Received 25 February 2025; Received in revised form 8 July 2025; Accepted 10 July 2025

Available online 12 July 2025

0959-8049/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

patients that had progressed after two previous lines of treatment (>90 %) [1]. Objective response rate (ORR) according to Response Evaluation Criteria In Solid Tumors (RECIST) version 1.1 [2] favored the FTD/TPI + bevacizumab arm (6.1 % and 1.2 % of patients in the FTD/TPI + bevacizumab and FTD/TPI treatment groups, respectively), which is a relatively modest improvement [1].

RECIST classifies partial response (PR) as tumor shrinkage (TS) of ≥ 30 % and progressive disease (PD) as tumor growth of ≥ 20 %, while any tumor growth or shrinkage between these two values is classified as stable disease (SD) [2]. In addition, RECIST does not consider timing, depth or duration of response. Thus, the population in SUNLIGHT with SD following treatment is heterogeneous, and includes patients with TS and patients with tumor growth outside of RECIST classifications of PD and PR [3].

It has been suggested that tumor evaluation using RECIST may not capture the full spectrum of response to treatment [4], and there is a need to explore different parameters to assess response to treatment in mCRC clinical trials, with a focus on dynamic changes of the target lesion. This is particularly relevant in the third line setting, in which ORR remains low and thus RECIST may be more equivocal than in earlier lines of treatment. Response-related parameters, such as early tumor shrinkage (ETS), duration of TS (DTS) and depth of response (DpR), have been shown to correlate with OS across a number of solid tumors, including mCRC [5–9].

Thus, the aim of this *post hoc* analysis of the SUNLIGHT trial was to assess response to treatment with FTD/TPI + bevacizumab and FTD/TPI using ETS, DTS and DpR and the association of these parameters with OS, PFS and global health status (GHS) from QLQ-C30 (Quality of Life Questionnaire Core 30, from European Organisation for Research and Treatment of Cancer).

2. Methods

2.1. Trial design and population

The design of the SUNLIGHT (NCT04737187) trial has been described previously [1,10]. Briefly, this study included patients aged ≥ 18 years old with histologically confirmed unresectable adenocarcinoma of the colon or rectum and had an Eastern Cooperative Oncology Group performance status (ECOG PS) of ≤ 1 . The inclusion criteria for SUNLIGHT stated that patients must have progressed on their previous line of treatment and received no more than two previous chemotherapy regimens (including a fluoropyrimidine, irinotecan, oxaliplatin, an anti-VEGF monoclonal antibody and an anti-EGFR monoclonal antibody for RAS wild-type patients) and had progressed, or were intolerant to, their last line of treatment.

Patients were randomly assigned to receive FTD/TPI (35 mg/m² twice a day on days 1–5 and 8–12 every 28-day cycle) or FTD/TPI in combination with bevacizumab (5 mg/kg on days 1 and 15).

2.2. Assessments

Tumor evaluations were conducted according to RECIST version 1.1 and involved clinical examination of image-based assessments of the chest, abdomen, and pelvis at a minimum (other localizations could be included if clinically indicated) [2]. Tumor assessments were performed at baseline and every 2 cycles (8 weeks) until radiologic disease progression. After progression, follow-up OS assessments continued every 8 weeks, until death or end of study. At baseline, images obtained before receiving a patient's written informed consent could be used if the date of the images was within 28 days of randomization, and if they were in line with the methods and techniques used for tumor evaluation in this study.

For the QLQ-C30 analyses, patients that completed at least one questionnaire at baseline and during the study period were analyzed in their respective treatment groups. Baseline values, post-baseline values

and changes from baseline in the GHS were collected at the beginning of each cycle of 28 days and at withdrawal visit. The QLQ-C30 consists of 30 questions and 15 scales including GHS quality of life (QoL) scale [11].

2.3. Endpoints

In this *post hoc* analysis, TS was defined as any decrease from baseline of the sum of the longest diameters of the target lesions. ETS was specified as TS occurring at the first tumor assessment (cycle 2 or before 8 weeks). DpR was defined as the maximum percentage change from baseline of the sum of the longest diameters of target lesions. DTS was defined as the time from first documented TS to the first documented increase in tumor size (from nadir), PD (clinical or radiological) or death, whichever came first.

2.4. Statistics

All analyses were performed in the tumor evaluable population (TUMEV), defined as all patients with measurable disease that had at least one evaluable baseline and post-baseline tumor assessment. Time to event endpoints were analyzed using Kaplan-Meier methodology and compared between treatment groups with a 2-sided log-rank test. Differences in TS/ETS rates and DpR between treatment groups were made using Fisher's exact test (two-sided) and two-sided Wilcoxon test, respectively.

A multivariate Cox proportional hazard regression analysis was performed to assess the relationship between ETS and OS/PFS. This analysis was adjusted for known prognostic factors (ECOG PS, location of primary disease, prior surgical resection, number of metastatic sites, neutrophil-to-lymphocyte ratio, prior bevacizumab) and stratification factors (geographic region, time since diagnosis of first metastasis, RAS mutational status). Statistical significance for factors were evaluated using Wald chi-square tests. An unstratified Cox regression model was used to assess the association between DpR and relative change from baseline at first tumor assessment, and both PFS and OS.

3. Results

Of 492 patients recruited in SUNLIGHT [1], 246 were randomized to each treatment group, and 227 and 218 were included in the TUMEV from the FTD/TPI + bevacizumab and FTD/TPI groups, respectively (Figure 1). In the FTD/TPI + bevacizumab group, 48 % and 39 % had TS and ETS, respectively (Figure 1). In the FTD/TPI group, 21 % and 17 % had TS and ETS, respectively (Figure 1). Baseline characteristics of patients in each treatment group, with and without TS and ETS are summarized in Table 1. No major differences in baseline characteristics were observed between the groups.

3.1. Tumor shrinkage and depth of response

The magnitude of DpR, TS and ETS was greater in the FTD/TPI + bevacizumab treatment group than the FTD/TPI treatment group. There was a 27.4 % difference in TS rate between treatment groups ($P < 0.0001$), and there was a 22.7 % difference in ETS rate between treatment groups ($P < 0.0001$). TS for each treatment group is summarized as a waterfall plot in Figure 2A.

In patients achieving TS, median DTS was 3.7 months in patients treated with FTD/TPI + bevacizumab and 2.2 months in patients treated with FTD/TPI (HR: 0.55 [95 % CI: 0.38, 0.78]; $P = 0.00082$; Figure 2B). In patients achieving ETS, median DTS was 3.8 months in patients treated with FTD/TPI + bevacizumab and 2.1 months in patients treated with FTD/TPI (HR: 0.34 [95 % CI: 0.22, 0.53]; $P < 0.0001$; Figure 2C). The magnitude of DpR was greater with FTD/TPI + bevacizumab treatment than with FTD/TPI treatment in the TUMEV and in those with TS and ETS (Supplementary Table 1).




Fig. 1. CONSORT diagram summarizing the populations used for analysis (original CONSORT diagram for the SUNLIGHT study published in Prager et al. [1]).

*Tumor shrinkage is defined as any decrease in sum of the longest diameter of the target lesion. **ETS is defined as tumor shrinkage that occurred before or at cycle 2 or 8 weeks. Percentages shown here are of the TUMEV. ETS, early tumor shrinkage. FTD/TPI, trifluridine/tipiracil. TUMEV, tumor evaluable population.

Table 1

Baseline characteristics and disease characteristics of patients with and without TS and ETS.

	With TS		Without TS		With ETS		Without ETS	
	FTD/TPI + bev (n = 110)	FTD/TPI (n = 46)	FTD/TPI + bev (n = 117)	FTD/TPI (n = 172)	FTD/TPI + bev (n = 88)	FTD/TPI (n = 35)	FTD/TPI + bev (n = 139)	FTD/TPI (n = 183)
Mean (SD) age, years	61.6 (10.0)	63.6 (10.4)	61.1 (12.4)	62.4 (11.2)	62.1 (9.2)	63.4 (11.2)	60.9 (12.4)	62.5 (11.0)
Gender, n (%)								
Female	52 (47.3)	21 (45.7)	62 (53.0)	81 (47.1)	41 (46.6)	15 (42.9)	73 (52.5)	87 (47.5)
Male	58 (52.7)	25 (54.4)	55 (47.0)	91 (52.9)	47 (53.4)	20 (57.1)	66 (47.5)	96 (52.5)
Ethnic origin, n (%)								
White	98 (94.2)	43 (95.6)	100 (95.2)	153 (96.2)	78 (94.0)	33 (97.1)	120 (95.2)	163 (95.9)
Black or African American	1 (1.0)	0 (0.0)	2 (1.9)	3 (1.9)	1 (1.2)	0 (0.0)	2 (1.6)	3 (1.8)
Other	5 (4.8)	2 (4.4)	3 (2.9)	3 (1.9)	4 (4.8)	1 (2.9)	4 (3.2)	4 (2.4)
ECOG PS, n (%)								
0	55 (50.0)	20 (43.5)	56 (47.9)	77 (44.8)	45 (51.1)	16 (45.7)	66 (47.5)	81 (44.3)
1	55 (50.0)	26 (56.5)	61 (52.1)	94 (54.7)	43 (48.9)	19 (54.3)	73 (52.5)	101 (55.2)
2	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.6)
Time since first metastasis, n (%)								
<18 months	45 (40.9)	14 (30.4)	50 (42.7)	80 (46.5)	37 (42.1)	12 (34.3)	58 (41.7)	82 (44.8)
≥ 18 months	65 (59.1)	32 (69.6)	67 (57.3)	92 (53.5)	51 (58.0)	23 (65.7)	81 (58.3)	101 (55.2)
Liver metastasis, n (%)	81 (73.6)	26 (56.5)	100 (85.5)	146 (84.9)	68 (77.3)	20 (57.1)	113 (81.3)	152 (83.1)
Number of metastatic sites, n (%)								
1–2	74 (67.3)	28 (60.9)	70 (59.8)	100 (58.1)	55 (62.5)	22 (62.9)	89 (64.0)	106 (57.9)
≥ 3	36 (32.7)	18 (39.1)	47 (40.2)	72 (41.9)	33 (37.5)	13 (37.1)	50 (36.0)	77 (42.1)
Location of primary disease, n (%)								
Right	25 (22.7)	12 (26.1)	34 (29.1)	54 (31.4)	19 (21.6)	8 (22.9)	40 (28.8)	58 (31.7)
Left	85 (77.3)	34 (73.9)	83 (70.9)	118 (68.6)	69 (78.4)	27 (77.1)	99 (71.2)	125 (68.3)
Prior surgical resection, n (%)	84 (76.4)	40 (87.0)	63 (53.9)	119 (69.2)	66 (75.0)	30 (85.7)	81 (58.3)	129 (70.5)
Prior bev, n (%)								
Yes	64 (58.2)	32 (69.6)	97 (82.9)	122 (70.9)	50 (56.8)	23 (65.7)	111 (79.9)	131 (71.6)
Prior anti-cancer agents, n (%)								
Fluoropyrimidine	110 (100.0)	46 (100.0)	117 (100.0)	172 (100.0)	88 (100.0)	35 (100.0)	139 (100.0)	183 (100.0)
Irinotecan	110 (100.0)	46 (100.0)	117 (100.0)	171 (99.4)	88 (100.0)	35 (100.0)	139 (100.0)	182 (99.5)
Oxaliplatin	107 (97.3)	45 (97.8)	117 (100.0)	170 (98.9)	87 (98.9)	35 (100.0)	137 (98.6)	180 (98.4)

Bev, bevacizumab. CI, confidence interval. ECOG PS, Eastern Cooperative Oncology Group performance status. ETS, early tumor shrinkage. FTD/TPI, trifluridine/tipiracil. HR, hazard ratio. OS, overall survival. PFS, progression-free survival. SD, standard deviation. TS, tumor shrinkage.




Fig. 2. Depth of response and tumor shrinkage according to treatment group for individual patients in the TUMEV (A), duration of response in patients receiving FTD/TPI + bevacizumab and FTD/TPI in the TUMEV with tumor shrinkage (B), and with early tumor shrinkage (C). Bev, bevacizumab. BOR, best overall response. CI, confidence interval. FTD/TPI, trifluridine/tipiracil. HR, hazard ratio. NE, non-evaluable. PD, progressive disease. PR, partial response. SD, stable disease. TS, tumor shrinkage. TUMEV, tumor evaluable population.

3.2. Association between survival, early tumor shrinkage and depth of response

In the TUMEV, median OS was longer in patients with ETS compared to patients without ETS (HR: 0.43 [95 % CI: 0.32, 0.57], $P < 0.0001$; Figure 3A). In both treatment groups, median OS was prolonged in patients achieving ETS compared to those not achieving ETS (HR: 0.39 [95 % CI: 0.27, 0.58], $P < 0.0001$ for the FTD/TPI + bevacizumab treatment group and HR: 0.61 [95 % CI: 0.40, 0.95], $P = 0.026$ for the FTD/TPI group; Figures 3B and 3C and Supplementary Figure 1A). Median OS was longer in the FTD/TPI + bevacizumab treatment group than in the FTD/TPI treatment group, regardless of whether ETS was achieved (HR: 0.75 [95 % CI: 0.58, 0.97], $P = 0.030$ for the subgroup without ETS and HR: 0.48 [0.28, 0.80], $P = 0.005$ for the subgroup with ETS; Supplementary Figure 1B).

A multivariate Cox proportional hazard regression analysis, adjusted for known prognostic factors, showed that the OS and PFS benefit in the FTD/TPI + bevacizumab group compared to the FTD/TPI group was maintained with and without ETS (Table 2). A Cox proportional hazard analysis confirmed that DpR and ETS were likely associated with both OS and PFS in the FTD/TPI + bevacizumab treatment group and the FTD/TPI treatment group (Supplementary Table 2).

3.3. Association between QoL deterioration and early tumor shrinkage

In patients receiving FTD/TPI + bevacizumab, GHS was maintained for longer in patients that achieved ETS compared to those without ETS (6.1 vs 4.4 months; $P = 0.031$; Supplementary Figures 2A and 3). Similarly, in patients receiving FTD/TPI, GHS was maintained for longer in patients with ETS compared to those without ETS (4.7 vs 3.2 months; $P = 0.16$; Supplementary Figures 2B and 3).

4. Discussion

In this *post hoc* analysis of SUNLIGHT, a higher proportion of TS and ETS and a greater magnitude of DpR was observed in patients receiving FTD/TPI + bevacizumab, than in patients receiving FTD/TPI. Additionally, the DTS was prolonged in the FTD/TPI + bevacizumab treatment group compared with the FTD/TPI group, in patients achieving TS and in patients achieving ETS. Therefore, this analysis suggests that the previously observed survival benefit of treatment with FTD/TPI + bevacizumab compared to FTD/TPI in SUNLIGHT is accompanied by improvement of the more refined metrics of tumor response (ETS and DpR).

Although, ORR based on RECIST was modest when compared to the improved PFS and OS magnitude demonstrated in SUNLIGHT [1], this additional analysis showed improved response to FTD/TPI + bevacizumab compared to FTD/TPI outside of the confines of the RECIST classifications (i.e., TS of $\geq 30\%$) [2]. The response to FTD/TPI + bevacizumab measured by ETS was likely associated with prolonged OS, which suggests that ETS in response to treatment may be associated with a survival benefit, despite not being a part of the RECIST classifications.

The use of ETS as a clinical endpoint has been widely investigated. Previous studies have shown that while ETS is a good prognostic factor, it may not be appropriate to use ETS as a surrogate marker for predicting treatment efficacy in mCRC [12]. However, in this analysis of patients in the third-line treatment setting, there was likely an association between OS and ETS, irrespective of treatment group, which is consistent with results of a previous meta-analysis of mCRC after first-line treatment [13]. While the use of ETS as an endpoint for clinical trials needs further exploration, its use in clinical practice could be beneficial. ETS may be a helpful addition to existing outcome measures as it may provide an earlier indication, from first tumor assessment, of potential future response [14]. In turn, this could optimize the management of patients with mCRC and aid clinical decision-making.

Furthermore, this analysis demonstrated an association between ETS

and prolonged time to GHS deterioration in both treatment groups, with a greater association in the FTD/TPI + bevacizumab treatment arm. This is of importance as some patients with advanced cancer report QoL to be as important as survival outcomes [15]. So, patients receiving FTD/TPI + bevacizumab that experience ETS are more likely to have an additional benefit of prolonged QoL.

DpR has been shown to be a valid surrogate endpoint for survival in mCRC patients receiving first-line cetuximab, panitumumab or bevacizumab-based chemotherapy and second-line cetuximab-based chemotherapy [16–19]. DpR has also been shown to be associated with OS and PFS in mCRC patients receiving first-line 5-fluorouracil, leucovorin, and irinotecan (FOLFIRI) + bevacizumab [6]. In this analysis, DpR was likely associated with both OS and PFS, and a greater magnitude of DpR was observed in patients treated with FTD/TPI + bevacizumab compared to those treated with FTD/TPI. Further work is needed to investigate the reliability of DpR as a predictor for overall response to treatment with FTD/TPI + bevacizumab.

A limitation of this analysis was that there was no central review of images from tumor assessments. There could be irregularities in the clinical examination of images between individual investigators, potentially introducing a level of bias. This was an unplanned *post hoc* analysis of SUNLIGHT and therefore conclusions should be taken with caution and used as the basis for hypothesis testing in future clinical trials investigating the use of FTD/TPI + bevacizumab. Another limitation of this study is the selected and small sample size for some analyses, particularly when investigating the association between OS and ETS. Future clinical trials with larger tumor evaluable populations should investigate the association between ETS and OS in patients treated with FTD/TPI + bevacizumab and put that in the perspective of common RECIST results.

5. Conclusion

In conclusion, this analysis of the SUNLIGHT trial showed that a higher proportion of patients receiving FTD/TPI + bevacizumab experienced TS and ETS, improved DpR and prolonged DTS than patients receiving FTD/TPI. ETS is associated with OS and deserves further evaluation in future prospective trials.

CRedit authorship contribution statement

Nadia Amellal: Writing – review & editing, Conceptualization. **Gerald W. Prager:** Writing – review & editing, Investigation, Conceptualization. **Julien Taieb:** Writing – review & editing, Investigation, Conceptualization. **Valentine Barboux:** Writing – review & editing, Formal analysis. **Lucas Roby:** Writing – review & editing, Writing – original draft. **Cristina Gravalos:** Writing – review & editing, Investigation. **Arinilda Campos Bragagnoli:** Writing – review & editing, Investigation. **Eric Van Cutsem:** Writing – review & editing, Investigation. **Elena Elez:** Writing – review & editing, Investigation. **Fortunato Ciardiello:** Writing – review & editing, Investigation. **Dominik P Modest:** Writing – review & editing, Investigation, Conceptualization. **Marwan Fakih:** Writing – review & editing, Investigation.

Funding

This analysis was funded by Institut de Recherches Internationales, Servier, France.

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests. Julien Taieb: Reports personal fees for speaker bureau and/or advisory role from Merck KGaA, Sanofi, Roche Genentech, MSD, BMS, Novartis, Astellas, Takeda, Ono Pharmaceutical, Boehringer-Ingelheim, Pierre




Fig. 3. Overall survival of patients with and without early tumor shrinkage in the TUMEV (A), in patients receiving FTD/TPI + bevacizumab (B) and in patients receiving FTD/TPI (C). Bev, bevacizumab. CI, confidence interval. ETS, early tumor shrinkage. FTD/TPI, trifluridine/tipiracil. HR, hazard ratio. NA, not applicable. TUMEV, tumor evaluable population.

Table 2
Association of ETS with OS and PFS in patients in the FTD/TPI + bevacizumab group versus patients in the FTD/TPI group.

Level	Factor	OS		PFS	
		HR [95 % CI]	P-value	HR [95 % CI]	P-value
FTD/TPI + bev versus FTD/ TPI ^a	With ETS	0.48 [0.27, 0.85]	0.013	0.26 [0.16, 0.43]	< 0.0001
	Without ETS	0.73 [0.56, 0.95]	0.021	0.47 [0.37, 0.60]	< 0.0001

^a HR calculated using a multivariate Cox proportional hazard regression model and corresponds to the treatment effect in each subgroup adjusted to all stratification factors (geographic region, time since diagnosis of first metastasis, RAS mutational status) and all significant prognostic factors (ECOG PS, location of primary disease, prior surgical resection, number of metastatic sites, neutrophil-to-lymphocyte ratio, prior bevacizumab). Bev, bevacizumab. CI, confidence interval. ETS, early tumor shrinkage. FTD/TPI, trifluridine/tipiracil. HR, hazard ratio. OS, overall survival. PFS, progression-free survival.

Fabre and Amgen, outside the submitted work and from Servier.

Dominik P Modest: Reports personal fees for speaker bureau and/or advisory role from Merck, KGaA, Sanofi, Amgen, Astra Zeneca, MSD, BMS, Incyte IKF, Takeda, Taiho, GSK, Pierre Fabre, Regeneron, Boehringer-Ingelheim.

Marwan Fakih: Consultancy/advisory for Amgen, Taiho Pharm., Bayer, Pfizer, Seagen, GSK, Incyte, Nouscom, Roche/Genentech, Mirati Ther. BMS, Eisai, and Merck. Speaker's bureau for Guardant Health; Honoraria, Amgen; Research Funding, Amgen, Verastem.

Fortunato Ciardiello: Consulting/advisory for Amgen, Merck KGaA, Pfizer, Roche/Genentech, Bayer US, LLC. Receives research funding from Amgen, BMS, Ipsen, Merck KGaA, Merck Sharp & Dohme, Roche/Genentech, Servier, Symphogen and Bayer US, LLC.

Eric Van Cutsem: Participation to advisory boards for Abbvie, Agenus, ALX, Amgen, Arcus Biosciences, Astellas, Astrazeneca, Bayer, Beigene, Bexon Clinical, Biontech, Boehringer Ingelheim, Bristol-Myers Squibb, Canfour, Daiichi, Debiopharma, Elmedix, Eisai, Galapagos, GSK, Hookipa Biotech, Incyte, Ipsen, Lilly, Merck Sharp & Dohme, Merck KGaA, Mirati, Novartis, Nordic, Pierre Fabre, Pfizer, Roche, Seattle Genetics, Servier, Simcere, Takeda, Taiho, and Terumo.

Elena Elez: Reports personal financial interest in Agenus, Amgen, Bayer, BMS, Boehringer Ingelheim, Cure Teq AG, GlaxoSmithKline, Hoffman La-Roche, Janssen, Lilly, Medscape, Merck Serono, MSD, Novartis, Organon, Pfizer, Pierre Fabre, Repare Therapeutics Inc., RIN Institute Inc., Sanofi, Seagen International GmbH, Servier, and Takeda

Cristina Gravalos: Reports personal fees for speaker bureau and/or advisory role from Servier, Merck, Amgen, Incyte and PierreFabre.

Arinilda Campos Bragagnoli: Reports speaker fees from Servier, Ipsen, BMS, Amgen, and Astrazeneca.

Valentine Barboux, Lucas Roby and Nadia Amellal: Employees of Servier.

Gerald W. Prager: Consulting/advisory for Amgen, Daiichi Sanyo Europe GmbH, Incyte, Merck Serono, Pierre Fabre, Roche/Genentech, Servier, Lilly O., Arcus Bioscience, and Bayer US LLC.

Acknowledgements

The authors thank Nicola Lloyd-Wood of ESP Medical Communications Ltd, Crowthorne, UK, for providing editorial support, which was sponsored by Servier, in accordance with Good Publication Practice

guidelines.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.ejca.2025.115644.

References

[1] Prager GW, Taieb J, Fakih M, Ciardiello F, Van Cutsem E, Elez E, et al. Trifluridine-tipiracil and bevacizumab in refractory metastatic colorectal cancer. *N Engl J Med* 2023;388:1657–67.

[2] Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228–47.

[3] Primavesi F, Fadinger N, Biggel S, Braunwarth E, Gasser E, Sprung S, et al. Early response evaluation during preoperative chemotherapy for colorectal liver metastases: combined size and morphology-based criteria predict pathological response and survival after resection. *J Surg Oncol* 2020;121:382–91.

[4] Venook AP, Tabernero J. Progression-free survival: helpful biomarker or clinically meaningless end point? *J Clin Oncol* 2015;33:4–6.

[5] Heinemann V, von Weikersthal LF, Decker T, Kiani A, Vehling-Kaiser U, Al-Batran SE, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2014;15:1065–75.

[6] Stintzing S, Modest DP, Rossius L, Lerch MM, von Weikersthal LF, Decker T, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab for metastatic colorectal cancer (FIRE-3): a post-hoc analysis of tumour dynamics in the final RAS wild-type subgroup of this randomised open-label phase 3 trial. *Lancet Oncol* 2016;17:1426–34.

[7] Piessevaux H, Buyse M, De Roock W, Prenen H, Schlichting M, Van Cutsem E, et al. Radiological tumor size decrease at week 6 is a potent predictor of outcome in chemorefractory metastatic colorectal cancer treated with cetuximab (BOND trial). *Ann Oncol* 2009;20:1375–82.

[8] Xie X, Li X, Yao W. A narrative review: depth of response as a predictor of the long-term outcomes for solid tumors. *Transl Cancer Res* 2021;10:1119–30.

[9] Heinemann V, Stintzing S, Modest DP, Giessen-Jung C, Michl M, Mansmann UR. Early tumour shrinkage (ETS) and depth of response (DpR) in the treatment of patients with metastatic colorectal cancer (mCRC). *Eur J Cancer* 2015;51:1927–36.

[10] Tabernero J, Taieb J, Prager GW, Ciardiello F, Fakih M, Leger C, et al. Trifluridine/tipiracil plus bevacizumab for third-line management of metastatic colorectal cancer: SUNLIGHT study design. *Future Oncol* 2021;17:1977–85.

[11] Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993;85:365–76.

[12] Petrelli F, Pietrantonio F, Cremolini C, Di Bartolomeo M, Coinu A, Lonati V, et al. Early tumour shrinkage as a prognostic factor and surrogate end-point in colorectal cancer: a systematic review and pooled-analysis. *Eur J Cancer* 2015;51:800–7.

[13] Colloca GA, Venturino A, Guarneri D. Early tumor shrinkage after first-line medical treatment of metastatic colorectal cancer: a meta-analysis. *Int J Clin Oncol* 2019;24:231–40.

[14] Douillard JY, Siena S, Peeters M, Koukakis R, Terwey JH, Tabernero J. Impact of early tumour shrinkage and resection on outcomes in patients with wild-type RAS metastatic colorectal cancer. *Eur J Cancer* 2015;51:1231–42.

[15] Meropol NJ, Egleston BL, Buzaglo JS, Benson AB, 3rd, Cegala DJ, et al. Cancer patient preferences for quality and length of life. *Cancer* 2008;113:3459–66.

[16] Van Cutsem E, Kohne CH, Hitre E, Zaluski J, Chang Chien CR, Makhson A, et al. Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer. *N Engl J Med* 2009;360:1408–17.

[17] Taieb J, Rivera F, Siena S, Karthaus M, Valladares-Ayerbes M, Gallego J, et al. Exploratory analyses assessing the impact of early tumour shrinkage and depth of response on survival outcomes in patients with RAS wild-type metastatic colorectal cancer receiving treatment in three randomised panitumumab trials. *J Cancer Res Clin Oncol* 2018;144:321–35.

[18] Cremolini C, Loupakis F, Antoniotti C, Lonardi S, Masi G, Salvatore L, et al. Early tumor shrinkage and depth of response predict long-term outcome in metastatic colorectal cancer patients treated with first-line chemotherapy plus bevacizumab: results from phase III TRIBE trial by the Gruppo Oncologico del Nord Ovest. *Ann Oncol* 2015;26:1188–94.

[19] Osumi H, Matsusaka S, Suenaga M, Shinozaki E, Mizunuma N. Associations between deepness of response and clinical outcomes among Japanese patients with metastatic colorectal cancer treated with second-line FOLFIRI plus cetuximab. *Oncol Targets Ther* 2015;8:2005–13.